

Care Bundles Recording Tool-Each shift

Peripherally Inserted Central Catheter

Use in conjunction with OLCHC (2013) Guidelines for Clinical Staff on Central Venous Access Devices & Careplan 9, 9a, 9b
 Tick ✓ each element of care bundle. To be 100% compliant all boxes must be ticked.

	Date:																			
	Time																			
Date PICC Inserted: ____ / ____ / 20__	Line Day																			
Date PICC Removed: ____ / ____ / 20__																				
Reason for line removal:																				
1. Assess the need for PICC on each shift* <i>if TPN in progress ? whether Enteral feeding/medications can commence Can oral medications be commenced</i> Please tick																				
2. Hand hygiene is performed before and after all line maintenance/access procedures. Please tick																				
3. Appropriate ANTT Level is used Please tick ANTT Level 2: accessing if 'breaking' the line ANTT Level 3: accessing via a needle free device																				
4. CVC site intact Please tick if site normal or X if not <i>Consider the following when observing the PICC site</i> a. Normal appearance b. Redness c. Redness, tenderness d. Redness, tenderness and discharge e. Other (temperature, bleeding)																				
5. Dressing replaced within 7 days or more often if required <i>(sterile transparent semi permeable dressing Veni-gard®/IV 3000). Change every 7 days)</i> Please tick																				
6. Disinfection solution is used to clean the insertion site during dressing changes. <i>(< 2 months corrected gestational age 0.5% Chlorohexidine in aqueous solution or older 2% Chlorohexidine in 70% alcohol)</i> Please tick																				
7. Change needle free device weekly Please tick <i>(Check date when needlefree change is due)</i>																				
8. Alcohol hub decontamination is performed before and after each hub access. <i>(Use OLCHC approved disinfection wipe)</i> Please tick																				
Initials																				
NMBI PIN																				

- At the beginning of each shift complete the care bundle above.
- *Under ' Assess the need for PICC ' if line not in use always place an x in the box and acknowledge the reason on the surveillance sheet
- If non compliant with bundle re-evaluation may be required following nursing care.
- Use a ✓ in the boxes provided for a yes. An X for no

Patient Name.....

HCR no.....

Ward

Care Bundles Recording Tool-Each shift

Peripherally Inserted Central Catheter Care Bundle

Use in conjunction with OLCHC (2013) Guidelines for Clinical Staff on Central Venous Access Devices & Careplan 9, 9a, 9b
 Tick ✓ each element of care bundle. To be 100% compliant all boxes must be ticked.

	Date:																						
	Time																						
Date PICC Inserted: ____ / ____ / 20__	Line																						
Date PICC Removed: ____ / ____ / 20__	Day																						
Reason for line removal:																							
1. Assess the need for PICC each shift ✖ <i>if TPN in progress ? whether Enteral feeding/medications can commence</i> <i>Can oral medications be commenced</i> <i>Please tick</i>																							
2. Hand hygiene is performed before and after all line maintenance/access procedures. <i>Please tick</i>																							
3. Appropriate ANTT Level is used <i>Please tick</i> ANTT Level 2: accessing if 'breaking' the line or TPN ANTT Level 3: accessing via a needle free device																							
4. PICC site intact <i>Please tick if site normal or X if not</i> <i>Consider the following when observing the PICC site</i> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr><td>a. Normal appearance</td></tr> <tr><td>b. Redness</td></tr> <tr><td>c. Redness, tenderness</td></tr> <tr><td>d. Redness, tenderness and discharge</td></tr> <tr><td>e. Other (temperature, bleeding)</td></tr> </table>	a. Normal appearance	b. Redness	c. Redness, tenderness	d. Redness, tenderness and discharge	e. Other (temperature, bleeding)																		
a. Normal appearance																							
b. Redness																							
c. Redness, tenderness																							
d. Redness, tenderness and discharge																							
e. Other (temperature, bleeding)																							
5. Dressing replaced within 7 days or more often if required <i>(sterile transparent semi permeable dressing Veni-gard®/IV 3000). Change every 7 days)</i> <i>Please tick</i>																							
6. Disinfection solution is used to clean the insertion site during dressing changes. <i>(< 2 months corrected gestational age 0.5% Chlorohexidine in aqueous solution or older 2% Chlorohexidine in 70% alcohol)</i> <i>Please tick</i>																							
7. Change needle free device weekly <i>Please tick</i> <i>(Check date when needlefree change is due)</i>																							
8. Alcohol hub decontamination is performed before and after each hub access. <i>(Use OLCHC approved disinfection wipe)</i> <i>Please tick</i>																							
Initials																							
NMBI PIN																							

- At the beginning of each shift complete the care bundle above.
- If non compliant with bundle re-evaluation may be required following nursing care.
- ✖ Under ' Assess the need for PICC ' if line not in use always place an x in the box and acknowledge the reason on the surveillance sheet.
- Use a ✓ in the boxes provided for a yes. An X for no

Patient Name.....

HCR no.....

Ward